

SPORT

The Scintillation Prediction Observations Research Task

James Spann¹, Charles Swenson², Otavio Durão³, Luis Loures⁴, Rod Heelis⁵, Rebecca Bishop⁶,
Guan Le⁷, Mangalathayil Abdu⁴, Linda Krause¹, Clezio Denardin³, Lidia Shibuya⁴

¹ NASA/MSFC, ² USU, ³ INPE, ⁴ ITA, ⁵ UTD, ⁶ Aerospace, ⁷ NASA/GSFC



SPORT

- **Joint United States / Brazil Science Mission Concept**

- **United States**


- Science Instruments

- **Brazil**

- Spacecraft


- Operations

Joint Science Data Analysis



Science

- The equatorial ionization anomalies



Bela Fejer, The Equatorial Ionosphere: A Tutorial
CEDAR Meeting, Seattle Washington, 2015

- Plasma Bubbles

GUVI (Same Local Time, Different Longitudes)

Why do bubbles form
and sometimes not at
Different Longitudes?



Kil, Hyosub, et al. "Coincident equatorial bubble detection by TIMED/GUVI and ROCSAT-1."
Geophysical research letters 31.3 (2004).



AEROSPACE




UtahStateUniversity



Plasma Bubbles



About 1.5 Hours to form a bubble



What is the state of the ionosphere here?

That leads to bubbles here ?



When bottom side seeding perturbations seem to always be present



Retterer, J. M., and P. Roddy. "Faith in a seed: on the origins of equatorial plasma bubbles." *Annales Geophysicae*. Vol. 32. No. 5. Copernicus GmbH, 2014.

Motion of Ionosphere (From Radar)

Morphology of the post-sunset vortex in the equatorial ionospheric plasma drift



Geophysical Research Letters

Volume 42, Issue 1, pages 9-14, 8 JAN 2015 DOI: 10.1002/2014GL062019

<http://onlinelibrary.wiley.com/doi/10.1002/2014GL062019/full#grl52441-fig-0001>



AEROSPACE

UTD

UAH
The University of Alabama in Huntsville




UtahStateUniversity



Neutral Winds and Conductivities

The importance of winds in different regions to triggering EPB particularly wind shears on the bottom of the ionosphere




[Electrodynamics of the equatorial evening ionosphere: 1. Importance of winds in different regions](#)

Authors A. D. Richmond, T.-W. Fang, A. Maute First Published: 7 March 2015 Vol: 120, Pages: 2118–2132 DOI: 10.1002/2014JA020934 <http://onlinelibrary.wiley.com/doi/10.1002/2014JA020934/full#jgra51625-fig-0001>



Bubbles Lead to Scintillations



David Hysell Altair Observations

Not all plasma bubble depletions are associated with scintillations?

Old Bubbles?

New Bubbles?



AEROSPACE



UAH


The University of Alabama in Huntsville



UtahStateUniversity

Science Goals

- 1) What is the state of the ionosphere that gives rise to the growth of plasma bubbles that extend into and above the F-peak at different longitudes?
- 2) How are plasma irregularities at satellite altitudes related to the radio scintillations observed passing through these regions?



AEROSPACE





UtahStateUniversity



Magnetic Field

Most ground/radar observations come from the American sector of unique magnetic geometry

IRGF 1960



IRGF 2010



AEROSPACE





UAH
The University of Alabama in Huntsville




UtahStateUniversity



Measurement and Instrumentation



Mission ConOps



Images: Charlie Gray



AEROSPACE





UAH
The University of Alabama in Huntsville



UtahStateUniversity




GPS Radio Occultation and Scintillation




SPORT Methodology


- The state of the ionosphere at early local times is related to the occurrence of scintillations at later local times.
 - How does this relation vary with longitude?
- Use case studies when SPORT ascending or descending node is within 17 to 24 LT sector.
- Examine ~15 degree longitude sectors



Methodology Strategy 1




Methodology Strategy 2




How often are ideal occultation

- Study using SPORT in ISS orbit.
- Over one orbit in the region within $\pm 30^\circ$
 - ~2 profiles over the previous orbit traces
 - ~2 profiles occur over successive orbit traces.





Conclusions

- CubeSat missions can be developed with a full/regular suite of science instruments.
- Mid inclination ISS orbits allow for the deconvolution of local time and longitude at low-latitudes
- A String of pearls mission to increase time resolution



SPORT Mission and ORBIT



Launch from ISS, 400 km Alt
~3 year life




SPORT Instruments


Ion Velocity Meter
UTD





GPS Occultation
Receiver
Aerospace



Langmuir, E-field,
Impedance Probe
USU




Fluxgate Magnetometer
NASA Goddard




SPORT Telemetry

Channel Name	Duty %	Rate Hz	Bit Rate bps	Alongtrack km
Ion Velocity Meter			1824	
Drifts	100%	2.00	288	3.83
Composition Sweeps	100%	2.00	1536	3.83
GPS RO			16000	
Dayside Tracking	50%	1.00	1000	7.66
Nightside Tracking	50%	50.00	15000	0.15
Langmuir Probe			1984	
DC Probe	100%	40.00	960	0.19
IV Sweeps	100%	0.04	491.52	191.43
Floating Probe Sweeps	100%	0.04	491.52	191.43
N _e Wave Power	100%	0.04	40.96	191.43
E-Field			1321	
DC field	100%	40.00	1280	0.19
E-Field Wave Power	100%	0.04	40.96	191.43
Impedance Probe			197	
I & Q Sweep	20%	0.04	196	191.43
Tracking	20%	40.00	192	0.19
Fluxgate Magnetometer			2880	
DC field	100%	40.00	2880	0.19
Star Imager			1500	
Star Subimage	100%	1.00	1500	7.66
Other			2624	
Science GPS timing	100%	40.00	2560	0.19
Science Housekeeping	100%	0.10	64	76.57
Rate collected on orbit			31210	



50 Mbit/second Downlink giving
a safety factor of 14



AEROSPACE



Utah State University

